Presentation to the Lake Roosevelt Water Quality Council

The Effects of Trace Elements on Water Quality and Biological Health in the Lake Roosevelt National Recreational Area: Columbia River In Cooperation with NPS



Objectives of Project

- Assess the movement of trace elements from the sediment to the overlying water column
- Assess the bioaccumulation of trace elements to the biota

 Assess the bioavailability and toxicity of the sediment to the biota




Process-Oriented Studies at 8 sites

- Release of metals from sediments
 - Analysis of ambient porewater
 (0 2 cm) from four replicate cores
 - Analysis of water overlying a 20-cm core over 4 weeks

- Bioavailability of metals
 - Two toxicity tests
 - One bioaccumulation study
 - Acid volatile sulfursimultaneously extracted metals
 - Sequential selective metals




Schematic of Sample Collection and Analysis





Total Metal Concentrations

 Analysis of solution by ICP-MS after digestion with nitric and hydrofluoric acid






Toxicity Test No. 1





Toxicity Test No. 2



Chironomus tentans (midge) survival and growth in laboratory tests



Bioaccumulation

Bioaccumulation Study





Simultaneously Extracted Metals (SEM) Normalized by Acid-Volatile Sulfide (AVS)

- EPA Method 376.3 –
- AVS measures the amount of labile sulfur
- SEM measures the amount of metals that are released with the labile sulfur
- Toxicity is rarely observed when the ratio of SEM:AVS is less than 1.0





Selective Extraction: Four Sequential Solutions

#1: Weakly Adsorbed: 0.11 M Acetic Acid Bond to Fe and Mn oxide coatings: 0.5 M Hydroxylamine Hydrochloride **Organic and sulfide mineral phases: #3:** 30% Peroxide stabilized with 1 M ammonium acetate Associated with lattice including slag: Nitric-hydrofluoric acid



Quality-Assurance—Pore and Incubation Waters for 72 Samples

- Four Filtering Blanks
- Two Standard Reference Material (SRM)
- One Incubation Column Blank
- One Incubation Replicate (5 samples)



QA—Total Metal Concentrations for 48 Sediment Solids

- Three laboratory Blanks
- Duplicates of Two SRMS (Four samples)
- Field Split (One Composite)



QA- Toxicity Tests for Each Site

- Two test species
 - Midge (Chironomus tentans)
 - Amphipod (Hyalella azteca)
- Using standard and approved test
 - ASTM 1706-00
 - Strict test performance criteria
 - Control and reference toxicants
- 3-4 Test Replicates per site
- Using established lab
 - Long term organism cultures and test performance history
 - Good Laboratory Practices (GLP) Documentation



QA- Bioaccumulation Test for Each Site

One species - Earthworm (Lumbriculus variegatus) (Survives very well in freshwater sediments)

- Using standard, peer reviewed and approved Test
 - ASTM E1688-97
 - Test has strict performance criteria
 - Control organism body burden measure before and after test
 - All test organism body burden measured after test
- 3-4 Test Replicates per test
- Using established lab
 - Long term organism cultures and test performance history
 - Follow GLP practices & documentation



QA—AVS-SEM for 8 HCI Solutions


- Matrix Blank (AVS & SEM)
- Processing Blanks (AVS & SEM)
- One Replicate (AVS & SEM)
- Calibrated on six elements, semiquantitative on all others (SEM)




QA—Sequential Selective Extraction for 32 Solutions in 4 Different Extracts

- Matrix Blanks (4 extracts)
- Processing Blanks (4 extracts)
- Two SRMs (2 replicates in 4 extracts)
- One Replicate (4 extracts)





Zn Concentrations at Proposed Sites: From Previous Studies





Water Science Center in Washington http://wa.water.usgs.gov/

